


# Purification method development for Liraglutide




Purification is the most critical step in the manufacturing process of peptide therapeutics. The right choice of chromatography media is crucial for cost-effective production. With its wide pH range (pH 2–10), YMC-Triart Prep C18-S provides you with full flexibility in the method development of peptide purification. Simple scale-up procedures ensure the

reproducible result at manufacture-scale. A method for the purification of liraglutide with high resolution (antidiabetic peptide therapeutic, marketed by Novo Nordisk as Victoza®) was successfully developed with YMC-Triart Prep C18-S under alkaline condition. The purity obtained for the target compound was 99.5%.


## Step 1 Column Screening




## Step 5 Influence of Buffer Concentration




## Step 2 Influence of Organic Solvent




## Step 6 Gradient Optimization




## Step 3 Influence of pH



## Result Purification Run



## Step 4 Influence of Buffer Type



## Theoretical Scaling Up Calculation

Column	YMC-Triart Prep C18-S (10 µm, 12 nm)		
Eluent	A) 20 mM HCOONH <sub>4</sub> -NH <sub>3</sub> (pH 8.5) B) Acetonitrile		
Detection	UV at 215 nm		
Temperature	Ambient		
Cycle time	60 min/run – 8 cycles/day		
Column dimension	250 x 100 mm ID	250 x 450 mm ID	250 x 600 mm ID
Flow rate	0.47 L/min	9.52 L/min	16.92 L/min
Loading/run	6.0 g	121.5 g	216.0 g
Fraction volume / run	1.4 L	28.6 L	50.8 L
Liraglutide recovery / run	2.6 g	53.4 g	94.9 g
Liraglutide recovery / day	20.8 g	427.2 g	759.2 g

